CBCS SCHEME

USN | | | | | | | | | | |

21MAT21

Second Semester B.E. Degree Examination, June/July 2023
Advanced Calculus and Numerical Methods

Time: 3 hrs. Max. Marks: 100

Note: Answer any FIVE full questions, choosing ONE full question from each module.

Important Note : 1. On completing your answers, compulsorily draw diagonal cross lines on the remaining blank pages.
2. Any revealing of identification, appeal to evaluator and /or equations written eg, 42+8 = 50, will be treated as malpractice.

Module-1

1 a. Evaluate $\int_{-c}^{c}\int_{-b}^{b}\int_{-a}^{a}(x+y+z)dxdydz$. **(06 Marks)**

b. Evaluate $\int_{0}^{1}\int_{x}^{\sqrt{x}} xydydx$ by changing the order of integration. **(07 Marks)**

c. Prove that $\pi(\frac{1}{2})=\sqrt{\pi}$, using definition of Gama function. **(07 Marks)**

OR

2 a. Evaluate $\int_{0}^{\infty}\int_{0}^{\infty} e^{-(x^2+y^2)}dxdy$ by changing into polar coordinates. **(06 Marks)**

b. Find the area between the parabolas y^2-4ax and x^2-4ay by using double integration. **(07 Marks)**

c. Show that $\beta(m,n)=\frac{\Pi(m)\Pi(n)}{\Pi(m+n)}$. **(07 Marks)**

Module-2

3 a. Find the directional derivative of $\phi = x^2yz+4xz^2$ at the point (1, –2, –1) along $2\hat{i}-\hat{j}-2\hat{k}$. **(06 Marks)**

b. If $\vec{F}=\nabla(xy^3z^2)$, find $\text{div}\vec{F}$ and $\text{curl}\vec{F}$ at the point (1, –1, 1). **(07 Marks)**

c. If $\vec{F}=(x+y+az)\hat{i}+(bx+2y-z)\hat{j}+(x+cy+2z)\hat{k}$, find a, b, c such that $\text{curl}\vec{F}=0$. **(07 Marks)**

OR

4 a. If $\vec{F}=xy\hat{i}+yz\hat{j}+zx\hat{k}$, evaluate $\int_C \vec{F}.d\vec{v}$ where 'c' is the curve represented by x = t, y = t^2, z = t^3, $-1\le t\le 1$. **(06 Marks)**

b. Using Green's theorem, evaluate $\int_C (xy+y^2)dx+x^2dy$, where 'c' is bounded by y = x and y = x^2. **(07 Marks)**

c. Apply Stoke's theorem to evaluate $\iint \text{curl}\vec{F}.\hat{n}ds$ where $\vec{F}=(x^2+y^2)\hat{i}-2xy\hat{j}$ taken around the rectangle bounded by the lines x = ±a, y = 0 and y = b. **(07 Marks)**

VTU-13-10-2023 01:06:18pm AD - AD - AD - AD - AD
13-10-2023 01:32:40pm AD - AD - AD - AD - AD

Module-3

5 a. Form a partial differential equation by eliminating arbitrary function from $Z = f(x+at) + g(x-at)$. **(06 Marks)**

b. Solve $\frac{\partial^2 z}{\partial x \partial y} = \sin x \sin y$ for which $\frac{\partial z}{\partial y} = -2\sin y$ when x = 0 and z = 0 when y is an odd multiple of $\frac{\pi}{2}$. **(07 Marks)**

c. Derive one dimensional heat equation. **(07 Marks)**

OR

6 a. Form a partial differential equation by eliminating arbitrary constant from $Z = (x-a)^2 + (y-b)^2$. **(06 Marks)**

b. Solve $(y-z)p + (z-x)q = x - y$. **(07 Marks)**

c. Solve $\frac{\partial^2 z}{\partial y^2} = z$ given that when y = 0, z = e^x and $\frac{\partial z}{\partial y} = e^x$. **(07 Marks)**

Module-4

7 a. The area of a circle (A) corresponding to diameter (D) is given below:

D	80	85	90	95	100
A	5026	5674	6362	7088	7854

Find the area corresponding to diameter 105 using an appropriate interpolation formula. **(06 Marks)**

b. Find a real root of $x^3 - 2x - 5 = 0$ using Regula-Falsi method correct to 3 decimal places whose root lies between 2 and 2.5. **(07 Marks)**

c. Evaluate $\int_0^{\pi/2} \sqrt{\cos\theta}\, d\theta$ by taking 7 ordinates by Simpson's 1/3rd rule. **(07 Marks)**

OR

8 a. Use Newton's divided difference formula to find f(4) given the data:

x	0	2	3	6
f(x)	–4	2	14	158

(06 Marks)

b. Use Newton-Raphson method to find a real root of $x\sin x + \cos x = 0$ near x = Π. Carry out the iterations upto 4 decimal places. **(07 Marks)**

c. Use Lagrange's interpolation formula to find y when x = 35 to the following data:

x	25	30	40	60
f(x)	50	55	70	95

(07 Marks)

VTU-13-10-2023 01:06:18pm AD - AD - AD - AD - AD - AD - AD
13-10-2023 01:32:40pm AD - AD - AD - AD - AD - AD - AD - AD

Module-5

9 a. Use the Taylor series method to find y(0.2) from $\frac{dy}{dx} = y + \sin x$, y(0) = 1. **(06 Marks)**

b. Use Runge-Kutta method of order 4, find y at x = 0.1, given that $\frac{dy}{dx} = 3e^x + 2y$, y(0) = 0 with h = 0.1. **(07 Marks)**

c. Apply Milne's predictor-corrector method, to find y(1.4) from $\frac{dy}{dx} = x^2 + \frac{y}{2}$ given that y(1) = 2, y(1.1) = 2.2156, y(1.2) = 2.4649, y(1.3) = 2.7514. **(07 Marks)**

OR

10 a. Use modified Euler's method to solve $\frac{dy}{dx} = x^2 + y$ with y(0) = 1, h = 0.05 at x = 0.1. **(06 Marks)**

b. Use Taylor series method to find y(0.1) from $\frac{dy}{dx} = x^2 + y^2$ with y(0) = 1. **(07 Marks)**

c. Use Runge-Kutta method of 4th order, find y(0.1) given that $\frac{dy}{dx} = 3x + \frac{y}{2}$, y(0) = 1 with h = 0.1. **(07 Marks)**

* * * * *

VTU-13-10-2023 01:06:18pm
AD - AD - AD - AD - AD - AD - AD - AD
13-10-2023 01:32:40pm
AD - AD - AD - AD - AD - AD - AD - AD